

Table S2. Characteristics of the reported cases with a mandibular metastasis from HCC

Variable	Value
Sex	
Male	63 (86.3)
Female	10 (13.7)
Unknown	4
Age (yr)	61 (16–88)
Symptoms ^{a)}	
Swelling	49 (72.1)
Pain	22 (32.4)
Abnormal sensation (paresthesia or numbness)	20 (29.4)
Bleeding	11 (16.2)
Mouth disturbance	5 (7.4)
Loosening of tooth	2 (2.9)
Unknown	9
Sites of mandibular metastasis ^{a)}	
Ramus	26 (40.0)
Body	23 (35.4)
Angle	18 (27.7)
Condyle	6 (9.2)
Unknown	12
Diagnosis of mandibular metastasis before confirmation of primary HCC	
Yes	45 (67.2)
No	22 (32.8)
Unknown	10
Local treatments for mandibular metastasis	
Radiotherapy	22 (48.9)
Surgical treatment ^{b)}	17 (37.8)
Surgical treatment ^{b)} followed by radiotherapy	4 (8.9)
No local treatment	2 (4.4)
Unknown	25

Values are presented as number (%) or median (range). Percentages were calculated excluding any cases with unknown values. HCC, hepatocellular carcinoma.

^{a)}Data summed to above 100% due to coexisting or overlapping conditions.

^{b)}Surgical treatment included curettage and tooth extraction.