

Table S2. Characteristic of the patients according to their treatments

	RT alone (n = 32)	Surgery alone (n = 17)	Surgery + RT (n = 12)	Chemotherapy ± surgery/RT (n = 8)	Total (n = 60)
Age (yr)					
<60	17 (53.1)	9 (52.9)	3 (25.0)	3 (37.5)	32 (46.4)
≥60	15 (46.9)	8 (47.1)	9 (75.0)	5 (62.5)	37 (53.6)
Tumor location					
Extramedullary	8 (25.0)	5 (29.4)	3 (25.0)	2 (25.0)	18 (26.1)
Bone	24 (75.0)	12 (70.6)	9 (75.0)	6 (75.0)	51 (73.9)
Tumor size (cm)					
<5	14 (43.8)	12 (70.6)	5 (41.7)	2 (25.0)	33 (47.8)
≥5	12 (37.5)	5 (29.4)	7 (58.3)	6 (75.0)	30 (43.5)
Unknown	6 (18.8)	0 (0.0)	0 (0.0)	0 (0.0)	6 (8.7)
Serum M protein present					
No	15 (46.9)	2 (11.8)	0 (0.0)	1 (12.5)	18 (26.1)
Yes	16 (50.0)	2 (11.8)	2 (15.4)	5 (62.5)	25 (36.2)
Unknown	1 (3.1)	13 (76.5)	10 (84.6)	2 (25.0)	26 (37.7)
Bence Jones proteinuria					
Absent	23 (71.9)	2 (11.8)	2 (15.4)	5 (62.5)	32 (46.4)
Present	5 (15.6)	0 (0.0)	0 (0.0)	1 (12.5)	6 (8.7)
Unknown	4 (12.5)	15 (88.2)	10 (84.6)	2 (25.0)	31 (44.9)
Serum free light chain ratio					
Normal	8 (25.0)	1 (5.9)	1 (8.3)	1 (12.5)	11 (15.9)
Abnormal	8 (25.0)	1 (5.9)	0 (0.0)	2 (25.0)	11 (15.9)
Unknown	16 (50.0)	15 (88.2)	11 (91.7)	5 (62.5)	47 (68.1)

Values are presented as number of patients (%).

RT, radiation therapy.