

# Role of radiotherapy for pancreatobiliary neuroendocrine tumors

Jeongshim Lee, MD, Jinhyun Choi, MD, Chihwan Choi, MD, Jinsil Seong, MD, PhD

Department of Radiation Oncology, Yonsei Cancer Center, Yonsei University College of Medicine, Seoul, Korea

**Purpose:** We investigated the role of radiotherapy (RT) for pancreatobiliary neuroendocrine tumors (PB-NETs).

**Materials and Methods:** We identified 9 patients with PB-NETs who received RT between January 2005 and March 2012. Of these 9 patients, 4 were diagnosed with NETs in the pancreas and 5 were diagnosed with NETs in the gallbladder. All patients received RT to the primary tumor or resection bed with a median total irradiation dose of 50.4 Gy, with or without chemotherapy.

**Results:** The tumor response rate and tumor control rate in the RT field were 60% and 100 %, respectively. All 4 patients who underwent surgery had no evidence of disease in the RT field. Of the 5 patients who received RT to the primary gross tumor, 1 had complete response, 2 had partial response, and 2 had stable disease in the RT field. The median time to progression was 11 months. Of the 9 patients, four patients had no progression, and 5 patients had progression of disease (locoregional, 2; distant, 2; locoregional/distant, 1). Of the 4 patients without progression, 3 were treated with RT in adjuvant or neoadjuvant setting, and one received RT to primary tumor. One patient experienced radiation-induced duodenitis at 3 months after concurrent chemoradiation without treatment-related mortality.

**Conclusion:** RT can yield local control for advanced PB-NETs. RT should be considered an essential part of multimodality treatment in management of advanced PB-NETs.

**Keywords:** Neuroendocrine tumors, Radiotherapy, Treatment outcome

## Introduction

Pancreatobiliary neuroendocrine tumors (PB-NETs) are uncommon malignancies arising from the neuroendocrine system of the pancreatobiliary tract. The annual incidence is estimated at <5 in 1,000,000 in Asian, United States, and European studies [1-5]. Improvement of diagnostic tools and growing awareness of gastrointestinal NETs have resulted in increased detection, and thus, increased incidence of PB-NETs worldwide. Halfdanarson et al. [6] reported an annual incidence of 2.2 in 1,000,000, covering a period of 27 years, with a higher

incidence in PB-NETs in recent decades. In Korea, Cho et al. [7] reported that the incidence of gastroenteropancreatic neuroendocrine tumors (GEP-NETs) has shown a remarkable increase during the last decade and the incidence of PB-NETs is 9.5% of all GEP-NETs.

For treatment of PB-NETs, a multidisciplinary approach including surgical resection and chemotherapy is acceptable [8], but advanced PB-NETs still remain a difficult therapeutic challenge because of their high malignant potential and resistance to conventional chemotherapy with etoposide, platinum agents, anthracyclines, streptozocin, and 5-fluorou-

Received 27 May 2013, Revised 11 June 2013, Accepted 15 July 2013.

Correspondence: Jinsil Seong, MD, PhD, Department of Radiation Oncology, Yonsei Cancer Center, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 120-749, Korea. Tel: +82-2-2228-8116, Fax: +82-2-312-9033, E-mail: JSSEONG@yuhs.ac

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

[www.e-roj.org](http://www.e-roj.org)

racil (5-FU) based agents [1,9]. Thus, there is not yet a consensus on appropriate therapeutic approaches in the management of PB-NETs. Radiotherapy (RT), or combined chemoradiation (CRT), has not been widely used in the management of PB-NETs, and also has not been incorporated into multidisciplinary management of PB-NETs.

In this study, we retrospectively analyzed clinical outcomes to investigate the potential role of RT for PB-NETs patients. This study provided a single institution experience of patients with PB-NETs who were treated with external beam radiation (EBRT) intended for curative purposes.

## Materials and Methods

### 1. Patient eligibility

We included patients diagnosed with PB-NETs who were treated with RT with or without chemotherapy at our institution between January 2005 and March 2012. We identified 9 patients (7 males, 2 females) who had a pathological diagnosis of PB-NETs and who received EBRT to the primary tumor or resection bed.

### 2. Radiotherapy technique

All patients received RT to the primary tumor or resection bed for a total irradiated dose of 45 Gy or above, with or without chemotherapy. All patients received 3-dimensional conformal radiotherapy with energies  $\geq 6$  MV generated by a linear accelerator or intensity modulated radiation therapy (IMRT) by tomotherapy HI-ART II (TomoTherapy Inc., Madison, WI, USA). The treatment volume consisted of the gross tumor volume, defined by pancreatic and locoregional radiographic abnormalities identified by contrast-enhanced computed tomography (CT), the clinical target volume, defined as the area at risk for subclinical microscopic disease, and the planning target volume, typically consisting of a 0.5-cm margin outside of the clinical target volume. The RT technique consisted of multi-field techniques. If the patients received RT to the primary tumor, after a dose of 41.4 Gy in 23 fractions, the field was reduced to protect the critical organ and then an additional boost of 5.4 to 9.0 Gy was delivered in fractions of 1.8 Gy.

### 3. Treatment outcome assessment

Patients were considered assessable if measurable disease on CT scan or positron emission tomography-CT was present prior to the initiation of treatment. Response to treatment in the RT

field was evaluated using revised Response Evaluation Criteria in Solid Tumors (ver. 1.1) criteria [10], with complete response (CR) defined as the disappearance of the all target lesions, partial response (PR) defined as at least a 30% decrease in the sum of diameters of target lesions and progressive disease (PD) defined as at least a 20% increase in the sum of diameters of target lesions. Stable disease (SD) was defined as neither sufficient shrinkage to qualify for PR nor sufficient increase to qualify for PD. Additionally, no evidence of disease (NED) was defined as no objective evidence of tumor recurrence for patients treated with surgery. The tumor response rate for primary tumor was the proportion of CR and PR, and tumor control rate for primary tumor or resection bed was the proportion of NED, CR, PR, and SD. Local failure was defined as tumor recurrence in the primary site, regional failure was tumor recurrence in the regional lymph nodes, and recurrences at other sites were regarded as distant failure. Survival was defined as the time from the date of initiation of RT to the date of death as a result of any cause. Time to progression (TTP) was defined as the time from the date of initiation of RT to the first objective documentation of tumor progression or to the time of death as a result of PD in the absence of previous documentation of objective PD.

### 4. Toxicity assessment

We defined acute toxicities as events occurring within 3 months from the start of RT. Late toxicities were defined as events occurring after 3 months from the start of RT. All toxicities were graded according to the Common Terminology Criteria for Adverse Events (ver. 3.0) and toxicities rated higher than grade 3 were considered severe.

## Results

### 1. Patient and treatment characteristics

The median age at the time of initiation of treatment was 54 years (range, 38 to 61 years). Seven patients (77.8%) were men. All patients had the Eastern Cooperative Oncology Group performance of 0 or 1. Of the 9 patients, 4 patients were diagnosed with NET in the pancreas and 5 patients were diagnosed with NET in the gallbladder. Prior to treatment, 7 patients had locally advanced NET, 1 patient had adjacent liver metastasis, and 1 patient had para-aortic lymph node and adjacent liver metastasis. Six patients were diagnosed with poorly differentiated NETs. Among 9 patients, 4 underwent surgical resection. One patient was treated with preoperative

RT followed by surgery and 3 patients were treated with surgery followed by postoperative RT. All four patients who underwent surgical resection had tumor resection with negative margins.

The median of the total irradiation dose was 50.4 Gy (range, 45.0 to 58.4 Gy). All patients completed the full course of RT with or without chemotherapy at the intended dose without dose reduction. Six patients received EBRT by a linear accelerator from 45.0 to 50.4 Gy at 1.8 Gy daily, 5 times a week, and 3 patients were treated with IMRT by tomotherapy from 55.7 to 58.4 Gy at 2.2 to 2.9 Gy daily, 5 times a week. Of the 9 patients, 7 received a combination of concurrent CRT, and 2 patients received RT alone. Four patients were treated with etoposide plus carboplatin, 2 patients were treated with 5-FU, and 1 patient was treated with etoposide, and carboplatin plus 5-FU with concurrent RT. Five patients were treated with CRT to primary tumor, and 4 patients underwent resection with adjuvant CRT/RT or neoadjuvant CRT. The patient and treatment characteristics are summarized in Table 1.

**2. Treatment outcome**


Of the 9 patients, four patients (45%) underwent surgery with NED in the RT field, despite absence of additional treatment. Of the remaining 5 patients, three patients (33%) who received concurrent CRT were diagnosed with locally advanced PB-NETs with or without distant metastasis and experienced CR or PR in the RT field. Fig. 1 illustrates the regression in tumor size of patient #5. Two (22%) of the 5 patients demonstrated SD in the RT field. The tumor response rate for primary tumor and tumor control rate for primary tumor or resection bed were 60% (20% CR, 40% PR, and 40% SD) and 100% (45% NED, 11% CR, 22% PR, and 22% SD), respectively.

The median follow-up period for surviving patients was 16 months (range, 13 to 59 months) and the median TTP was 11 months (range, 2 to 59 months). Four patients were alive without progression, 1 patient was alive with progression, and 4 patients were dead with progression. Progression of disease occurred in 5 patients with 2 locoregional failures, 2 distant failures, and 1 synchronous locoregional and distant failure. Three of 4 patients underwent surgery and one of 5 patients who received concurrent CRT were alive without disease. One patient (patient #6) who survived 5 years from preoperative concurrent CRT followed by surgery without tumor recurrence was diagnosed initially with locally advanced well differentiated NET without lymph node metastasis in the pancreas. The prominent pattern of mortality in patients was

**Table 1. Patient and treatment characteristics**

No.	Sex/age	Location	T	N	M	Grade	Remark	Tx modality	CTx	Total dose (Gy)	Fraction dose (Gy)	RT modality
1	M/53	Pancreas	T2	N1	M0	NA	NA	CRT	EP	50.4	1.8	3D CRT
2	F/61	GB	T3	N2	M0	PD	LA with liver invasion	CRT	EP	50.4	1.8	3D CRT
3	M/56	GB	T3	N1	M0	PD	LA with liver invasion	CRT	5-FU	58.4	2.5	IMRT
4	M/56	Pancreas	T2	N0	M1	NA	Adjacent liver metastasis	CRT	EP	55.9	2.2	IMRT
5	M/50	GB	T3	N2	M1	PD	LA with liver invasion, adjacent liver metastasis	CRT	EP	50.4	1.8	3D CRT
6	M/57	Pancreas	T2/3	N0	M0	WD	RM (-)	CRT → Op	EPF	55.7	2.9	IMRT
7	F/38	GB	T3	N1	M0	PD	RM (-)	Op → CRT	5-FU	45.0	1.8	3D CRT
8	M/41	GB	T2	N1	M0	PD	RM (-)	Op → RT	No	45.0	1.8	3D CRT
9	M/41	Pancreas	T3	N0	M0	MD	RM (-), LV/PNI (+/+)	Op → RT	No	50.4	1.8	3D CRT

T, tumor; N, node; M, metastasis; Tx, treatment; CTx, chemotherapy; RT, radiotherapy; NA, not applicable; GB, gallbladder; LA, locally advanced; EP, etoposide and carboplatin; RM, resection margin; 5-FU, 5-fluorouracil; CRT, chemoradiation; Op, operation; IMRT, intensity modulated radiation therapy; LV, lymphovascular invasion; PNI, perineural invasion.



**Fig. 1.** Abdominal computed tomography scans of 51-year-old male patient diagnosed as having gallbladder neuroendocrine tumor with lymph node involvement and adjacent liver metastasis (arrow) treated with radiotherapy (RT) to gross mass: (A) 1 month before RT, (B) 1 month after RT, and (C) 12 months after RT.

**Table 2.** Treatment outcomes of radiotherapy for pancreatobiliary neuroendocrine tumors

No.	Infield outcome	Recurrence	TTP (mo)	Remark	Last follow-up status	OS (mo)
1	SD	DM	2	SD of primary tumor and LN site, but developed liver and lung metastasis	DWD	28
2	SD	R	11	SD for 11 months, and then regional LN progression	AWD	15
3	PR	L + R + DM	5	PR for 5 months, and then primary tumor, LN and liver progression	DWD	8
4	PR	DM	3	PR of primary tumor site, but progressive liver metastasis	DWD	2
5	CR	No	-	(CR at 12 mo)	NED	12
6	NED	No	-	(NED for 5 yr)	NED	60
7	NED	No	-	(NED for 16 mo)	NED	16
8	NED	R	5	NED for 4 mo, and then regional LN recurrence	DWD	12
9	NED	No	-	(NED for 15 mo)	NED	15

TTP, time to progression; OS, overall survival; SD, stable disease; DM, distant metastasis; LN, lymph node; CR, complete response; PR, partial remission; PD, progressive disease; NED, no evidence of disease; L, local recurrence; R, regional recurrence; DWD, died with disease; AWD, alive with disease.

**Table 3.** Overview of published results

Author	No. of patients	Treatment modality	Remark
Strosberg et al. [11]	6	CCRT	Tumor response rate 80%
Torrise et al. [12]	3	RT	One patient: PR for 4 yr, another: PR for 7 mo
Rich [13]	3	RT	One patient: resolved local symptoms for 39 mo
Saif et al. [14]	6	CCRT or sequential CRT	Tumor response rate 100%; one patient: alive for 9 yr, another: alive for 4.5 yr
Zagar et al. [15]	17	CCRT	Local control rate 85%; 2-yr distant metastatic-free survival 46%
Present	9	CCRT or RT	Tumor response rate 60%; tumor control rate 100%; median TTP 11 mo; one patient: alive for 5 yr

CCRT, concurrent chemoradiation; RT, radiotherapy; CRT, chemoradiation; TTP, time to progression.

not locoregional failure, but distant metastasis. The treatment outcomes with survival status are shown in Table 2.

**3. Toxicity**

The three patients experienced acute hematologic toxicities

above grade 3. Two patients had grade 3 neutropenia and 1 had grade 3 thrombocytopenia during concurrent CRT. No acute non-hematologic grade 3 and 4 toxicities were observed. Late toxicity was observed in one patient who experienced grade 3 radiation-induced duodenitis at 3 months after

concurrent CRT. There were no treatment-related deaths.

## Discussion and Conclusion

RT has not historically played a major role in the treatment of PB-NETs. This study included a series of patients with PB-NETs treated with curative-intent RT for primary gross disease, or after resection. Our findings show that RT could be an optional treatment modality for achieving local control in management of PB-NETs. RT produced excellent tumor control rate and yielded local control in patients with advanced PB-NETs. In terms of toxicity associated with this treatment, 3 patients had acute hematologic grade 3 toxicities and 1 patient had late gastrointestinal grade 3 toxicities. However, there were no treatment-related deaths.

A review of the literature showed that the role of RT in the management for PB-NETs had not been extensively evaluated because of the low incidence of this tumor. Few reports have described RT use for PB-NETs. Strosberg et al. [11] reported a study in which 6 patients were treated with a combination of concurrent and sequential CRT for treatment of locally advanced pancreatic NETs. The radiologic response rate was 80% of patients and all patients tolerated the CRT. Torrisi et al. [12] reported 2 patients with locally advanced pancreatic NETs, 1 of whom was treated with a combination of intraoperative iodine-125 brachytherapy followed by 41.4 Gy EBRT, and the other patient was treated with a combination of EBRT to a dose of 45 Gy and chemotherapy. The first patient achieved a prolonged PR of over 4 years, whereas the second patient achieved a PR lasting 7 months and then had recurrence. Rich [13] described a patient with pancreatic NET who received 46.5 Gy EBRT and had resolved local symptoms for 39 months until the local tumor recurred. The patient died 61 months from the start of irradiation, despite subsequent chemotherapy consisting of 5-FU and streptozocin. Recently, Saif et al. [14] reported that all 6 patients with pancreatic NET treated with concurrent CRT consisting of capecitabine or 5-FU experienced clinical benefit and PR. One patient survived 9 years from treatment and another patient was alive at 4.5 years. These studies documenting clinical outcomes from RT in patients with PB-NETs have suggested that PB-NETs are not radioresistant, but are radioresponsive [11-15].

In our series, both locoregional failure and distant failure in patients receiving CRT/RT in adjuvant or neoadjuvant setting were higher than those who received CRT/RT to primary gross disease. In the adjuvant or neoadjuvant setting,

patients of PB-NETs treated with curative-intent RT had 25% locoregional failure and no distant failure. And patients who received concurrent CRT to the primary gross tumor had 40% locoregional failure and 60% distant failure. Zagar et al. [15] from the Duke University reviewed the 33 resected pancreatic NETs patients treated with or without RT in the adjuvant or neoadjuvant setting. Of the 33 patients, sixteen patients were treated with surgical resection alone, while 17 underwent resection with adjuvant or neoadjuvant RT. The 2-year local control rate was 85% for surgery with RT group and 90% for the surgery alone group ( $p = 0.38$ ). The authors concluded that the role of RT in the adjuvant management of pancreatic NETs remains unclear. However, no statistically significant difference of local control seen between both groups could be explained by factor that the potential probability of surgery with RT group had more patients with an aggressive tumor biology/more locally aggressive disease patients compared with the surgery group. Solorzano et al. [16] reported that, of patients with pancreatic NETs who undergo surgical resection with curative intent, 50% will develop a recurrence within 5 years. In present study, the finding that RT in the adjuvant setting had a 75% locoregional control rate without distant recurrence is useful information in the treatment of this disease. A summary of published studies with PB-NETs patients treated with RT are shown in Table 3.

There have been several studies that used a targeted peptide receptor analog in the management of advanced or metastatic NETs to improve clinical responses. Kwekkeboom et al. [17] reported the GEP-NETs showed high response rates to the somatostatin analog octreotate. The recent success of the biologic targeted agents, such as everolimus, sunitinib and endostatin, has renewed excitement about treating advanced pancreatic NETs [18-20]. Biologic targeted agents have anti-angiogenic properties, whereas RT is known to block the growth of tumor vasculature. So, the combination of targeted agents and EBRT might be an effective treatment for this disease based on the rationale that PB-NETs are highly vascular tumors.

Our study was limited because the analysis was retrospective and the number of patients was small, but there are very few studies on the treatment outcome of RT for PB-NETs because of the low incidence of this tumor and the low therapeutic application of RT for this disease. For all those reasons, prognostic parameter related progression and survival were unable to analysis in this study.

In conclusion, our experience provide that RT with or without

chemotherapy can be used as an effective treatment modality in the management of advanced PB-NETs. RT may induce tumor response/control and achieve local control of PB-NETs. The future role of RT in combination with chemotherapy or biologic modifiers for the treatment of PB-NETs remains to be established. We suggest that RT should be considered as a part of multimodality therapy in curative management of advanced PB-NETs.

### Conflict of Interest

No potential conflict of interest relevant to this article was reported.

### References

- Buchanan KD, Johnston CF, O'Hare MM, et al. Neuroendocrine tumors: a European view. *Am J Med* 1986;81:14-22.
- Lam KY, Lo CY. Pancreatic endocrine tumour: a 22-year clinicopathological experience with morphological, immunohistochemical observation and a review of the literature. *Eur J Surg Oncol* 1997;23:36-42.
- Garcia-Carbonero R, Capdevila J, Crespo-Herrero G, et al. Incidence, patterns of care and prognostic factors for outcome of gastroenteropancreatic neuroendocrine tumors (GEP-NETs): results from the National Cancer Registry of Spain (RGETNE). *Ann Oncol* 2010;21:1794-803.
- Hemminki K, Li X. Incidence trends and risk factors of carcinoid tumors: a nationwide epidemiologic study from Sweden. *Cancer* 2001;92:2204-10.
- Lawrence B, Gustafsson BI, Chan A, Svejda B, Kidd M, Modlin IM. The epidemiology of gastroenteropancreatic neuroendocrine tumors. *Endocrinol Metab Clin North Am* 2011;40:1-18.
- Halfdanarson TR, Rabe KG, Rubin J, Petersen GM. Pancreatic neuroendocrine tumors (PNETs): incidence, prognosis and recent trend toward improved survival. *Ann Oncol* 2008;19:1727-33.
- Cho MY, Kim JM, Sohn JH, et al. Current trends of the incidence and pathological diagnosis of gastroenteropancreatic neuroendocrine tumors (GEP-NETs) in Korea 2000-2009: multicenter study. *Cancer Res Treat* 2012;44:157-65.
- Kulke MH, Benson AB 3rd, Bergsland E, et al. Neuroendocrine tumors. *J Natl Compr Canc Netw* 2012;10:724-64.
- Moertel CG, Lefkopoulo M, Lipsitz S, Hahn RG, Klaassen D. Streptozocin-doxorubicin, streptozocin-fluorouracil or chlorozotocin in the treatment of advanced islet-cell carcinoma. *N Engl J Med* 1992;326:519-23.
- Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009;45:228-47.
- Strosberg J, Hoffe S, Gardner N, Choi J, Kvols L. Effective treatment of locally advanced endocrine tumors of the pancreas with chemoradiotherapy. *Neuroendocrinology* 2007;85:216-20.
- Torrizi JR, Treat J, Zeman R, Dritschilo A. Radiotherapy in the management of pancreatic islet cell tumors. *Cancer* 1987;60:1226-31.
- Rich TA. Radiation therapy for pancreatic cancer: eleven year experience at the JCRT. *Int J Radiat Oncol Biol Phys* 1985;11:759-63.
- Saif MW, Ng J, Chang B, Russo S. Is there a role of radiotherapy in the management of pancreatic neuroendocrine tumors (PNET)? *JOP* 2012;13:174-6.
- Zagar TM, White RR, Willett CG, et al. Resected pancreatic neuroendocrine tumors: patterns of failure and disease-related outcomes with or without radiotherapy. *Int J Radiat Oncol Biol Phys* 2012;83:1126-31.
- Solorzano CC, Lee JE, Pisters PW, et al. Nonfunctioning islet cell carcinoma of the pancreas: survival results in a contemporary series of 163 patients. *Surgery* 2001;130:1078-85.
- Kwekkeboom DJ, Teunissen JJ, Bakker WH, et al. Radiolabeled somatostatin analog [177Lu-DOTA0,Tyr3]octreotate in patients with endocrine gastroenteropancreatic tumors. *J Clin Oncol* 2005;23:2754-62.
- Yao JC, Shah MH, Ito T, et al. Everolimus for advanced pancreatic neuroendocrine tumors. *N Engl J Med* 2011;364:514-23.
- Raymond E, Dahan L, Raoul JL, et al. Sunitinib malate for the treatment of pancreatic neuroendocrine tumors. *N Engl J Med* 2011;364:501-13.
- Kulke MH, Bergsland EK, Ryan DP, et al. Phase II study of recombinant human endostatin in patients with advanced neuroendocrine tumors. *J Clin Oncol* 2006;24:3555-61.